

TYVASO® is approved for the treatment of pulmonary arterial hypertension (PAH; WHO Group 1) to improve exercise ability.

LIFE'S LITTLE MOMENTS.

BREATHE THEM IN WITH TYVASO

What you need to know
about TYVASO for PAH

TYVASO DELIVERS



Important Safety Information for TYVASO

Before you take TYVASO, tell your healthcare provider about all of your medical conditions, including if you:

- Have lung disease, such as asthma or chronic obstructive pulmonary disease (COPD)
- Have a lung infection

Please see Important Safety Information throughout and on page 11 and accompanying Full Prescribing Information in pocket.


WHO=World Health Organization.

TYVASO[®]
(treprostinil) INHALATION
SOLUTION

About your PAH treatment

People with pulmonary arterial hypertension (PAH) have an imbalance of 3 natural substances, causing the blood vessels in their lungs to become narrowed. When these 3 substances are out of balance, it is harder to keep blood flowing through the lungs and this causes symptoms of PAH.

There are medicines that treat each of those 3 imbalances, or pathways, associated with PAH. Each PAH medicine targets only one of those pathways.



What is prostacyclin? [pra-stuh·si·klin]

Prostacyclin is one of the pathways associated with PAH. Prostacyclin helps ensure blood vessels are healthy and stay open. People who have PAH may not produce enough prostacyclin in their lungs.

TYVASO helps to replace the natural prostacyclin you may be missing

TYVASO mimics some of the effects of natural prostacyclin, helping to open up blood vessels and keep the blood flowing through your lungs.


What is TYVASO?

TYVASO (treprostinil) is a prescription medicine used in adults to treat pulmonary arterial hypertension (PAH; WHO Group I), which is high blood pressure in the arteries of your lungs. TYVASO can improve the ability to exercise. Your ability to exercise decreases 4 hours after taking TYVASO. It is not known if TYVASO is safe and effective in children under 18 years of age.

Breathe it in...

TYVASO is inhaled directly to the lungs

TYVASO quickly delivers medicine to the site of disease, where it's needed



Important Safety Information for TYVASO

Before you take TYVASO, tell your healthcare provider about all of your medical conditions, including if you (continued):

- Have liver or kidney problems
- Have low blood pressure
- Have bleeding problems
- Are pregnant or plan to become pregnant. It is not known if TYVASO will harm your unborn baby.

Please see Important Safety Information throughout and on page 11 and accompanying Full Prescribing Information in pocket.



TYVASO may help you do what matters most

Walk farther in 6 minutes

In a clinical study, people who added TYVASO to their current PAH medicine walked farther—improving their 6MWD* in as soon as **6 weeks**.†

In a 6-minute walk test, people were able to walk:



Do more in your day

When you add TYVASO to your current treatment, you may be able to do things like:



*6-minute walk distance (6MWD) is a test to see how far you can walk in 6 minutes.

†In the 12-week clinical study of TYVASO, people taking an oral PAH medicine (bosentan or sildenafil) completed a 6-minute walk test at the beginning and end of the study.

Important Safety Information for TYVASO

Before you take TYVASO, tell your healthcare provider about all of your medical conditions, including if you (continued):

- Are breast-feeding or plan to breast-feed. It is not known if TYVASO passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with TYVASO.


Continuing the study

After the 12-week clinical study was over, patients who received TYVASO could continue taking it, and patients who had initially received placebo could start on TYVASO.

- *Nearly all of the patients who completed the initial study (97%) decided to continue as part of the long-term follow-up study.*

During that study, researchers followed patients to understand the dosing and safety of TYVASO. They also observed changes in their 6MWD, their ability to be active, and their rate of survival.

Patients who took TYVASO in the clinical study and continued to be observed...



It's important to note that in this study, all of the patients were taking TYVASO, and no comparison was made to people who were not taking TYVASO. So, it's hard to determine for sure if these results are because of TYVASO.

Important Safety Information for TYVASO

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TYVASO and other medicines may affect each other.

Please see Important Safety Information throughout and on page 11 and accompanying Full Prescribing Information in pocket.





TYVASO is delivered through a portable inhalation system.

Be sure to take TYVASO 4x daily as directed by your doctor so you get the full benefit of the medicine.



How to take TYVASO

Getting started and reaching your target dose

Most people start with 3 breaths, 4x daily and increase the number of breaths every week to a target dose of 9 breaths, 4x daily.



It took patients in the clinical study an average of 3 weeks to reach their target dose.

Reaching your target dose is important so you'll feel the most benefit from the medicine. Your doctor will work with you to increase the breaths per treatment session until you reach your target dose, to help you feel better sooner.

Important Safety Information for TYVASO

Especially tell your healthcare provider if you take:

- Medicines used to treat high blood pressure or heart disease
- Medicines that decrease blood clotting (anticoagulants)
- Water pills (diuretics)
- Gemfibrozil (Lopid) or rifampin (Rimactane, Rifadin, Rifamate, Rifater)

You may have side effects...

Don't let side effects stop you from getting the benefits of your medicine. **It's important to talk to your healthcare provider and continue taking TYVASO unless told to stop.** Be sure to contact your nurse about things you can do to help manage side effects while continuing treatment, such as:

COUGH/THROAT IRRITATION

- ✓ Be sure to keep the device level and take normal breaths during treatment, and avoid holding your breath once you inhale
- ✓ If needed, use a throat spray to numb your throat before treatment

NAUSEA

- ✓ Rinse your mouth with water after treatment

HEADACHE

- ✓ Use an over-the-counter pain reliever, like acetaminophen; talk to your doctor to see if slowing the pace at which you increase your dose could help

These tips may help you manage common side effects with TYVASO, based on healthcare provider experience. You should not think of this as medical advice.

Important Safety Information for TYVASO

What are the possible side effects of TYVASO?

TYVASO can cause **serious side effects**, including:

- **Low blood pressure** (symptomatic hypotension). If you have low blood pressure, TYVASO may lower your blood pressure more.
- **Bleeding problems.** TYVASO may increase the risk of bleeding, especially in people who take blood thinners (anticoagulants).

Please see Important Safety Information throughout and on page 11 and accompanying Full Prescribing Information in pocket.



TYVASO can go wherever your day takes you

You can take TYVASO with you throughout your day. The portable TYVASO device allows for dosing at home or on-the-go.

- Set up device **once a day**
- Each treatment session takes about **2 to 3 minutes**
- Plan for treatment sessions approximately every **4 waking hours**
- Clean device **once a night**



Important Safety Information for TYVASO

What are the possible side effects of TYVASO? (continued)

The **most common side effects** of TYVASO are cough, headache, throat irritation and pain, nausea, reddening of the face and neck (flushing), fainting or loss of consciousness, dizziness, and diarrhea. These are not all the possible side effects of TYVASO. Call your doctor for medical advice about side effects.

Please see Important Safety Information throughout and on page 11 and accompanying Full Prescribing Information in pocket.



Join the thousands who have chosen treatment with TYVASO

The decision to add TYVASO into your life is personal, but when you and your healthcare provider do, you should know that you are among thousands of people who have made the same decision. In fact, TYVASO is the most prescribed inhaled treatment for PAH.

Committed to helping people with PAH

TYVASO is produced by United Therapeutics Corporation, a company founded in 1996 by a couple whose daughter has PAH. At United Therapeutics, we are committed to helping people affected by PAH.

Important Safety Information for TYVASO

Before you take TYVASO, tell your healthcare provider about all of your medical conditions, including if you:

- Have lung disease, such as asthma or chronic obstructive pulmonary disease (COPD)
- Have a lung infection
- Have liver or kidney problems
- Have low blood pressure
- Have bleeding problems



Supporting you in many ways

It starts with a call from a Specialty Pharmacy

TYVASO is available only through Specialty Pharmacies, which focus on providing you with in-home training and 24-hour support. Your Specialty Pharmacy (SP) provider is part of your healthcare team.



What happens next: Your SP provider will call you.

Answer the call! When they first reach out to you, the SP provider may call you from an 866 or 877 number that you may not recognize. It's important to answer the call because they may not be able to leave a message for you due to privacy rules. Since they cannot get the medicine to you unless they speak to you, be sure to answer the call to get started and to stay on treatment.



An SP nurse will give you personal instructions on using TYVASO and provide ongoing support. Your SP nurse may visit or call regularly as you are getting comfortable using TYVASO.



Pay as little as \$5 per prescription

With the Co-Pay Assistance Card, eligible patients taking TYVASO who enroll may pay as little as a \$5 co-pay per month and may receive up to \$8000 in savings per year.

Learn how to enroll at www.UTcopay.com or call ASSIST at 1-877-864-8437.



Eligibility requirements for this Program are:

- You must be 18 years or older to use this Program
- Patients using Medicare, Medicaid, or any other state or federal government program to pay for their medications are not eligible. Patients who start utilizing government coverage during the term of the Program will no longer be eligible
- The Program is valid only for patients with commercial (also known as private) insurance who are taking the medication for an FDA-approved indication
- Eligible patients must be a resident of the United States or Puerto Rico. The Program is subject to additional state law restrictions. Patients residing in select states may not be eligible for the Program

Visit www.UTcopay.com for full Program details and Terms and Conditions.

Important Safety Information

Before you take TYVASO, tell your healthcare provider about all of your medical conditions, including if you:

- Have lung disease, such as asthma or chronic obstructive pulmonary disease (COPD)
- Have a lung infection
- Have liver or kidney problems
- Have low blood pressure
- Have bleeding problems
- Are pregnant or plan to become pregnant. It is not known if TYVASO will harm your unborn baby.
- Are breast-feeding or plan to breast-feed. It is not known if TYVASO passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with TYVASO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TYVASO and other medicines may affect each other.

Especially tell your healthcare provider if you take:

- Medicines used to treat high blood pressure or heart disease
- Medicines that decrease blood clotting (anticoagulants)
- Water pills (diuretics)
- Gemfibrozil (Lopid) or rifampin (Rimactane, Rifadin, Rifamate, Rifater)

What are the possible side effects of TYVASO?

TYVASO can cause **serious side effects**, including:

- **Low blood pressure** (symptomatic hypotension). If you have low blood pressure, TYVASO may lower your blood pressure more.
- **Bleeding problems.** TYVASO may increase the risk of bleeding, especially in people who take blood thinners (anticoagulants).

The **most common side effects** of TYVASO are cough, headache, throat irritation and pain, nausea, reddening of the face and neck (flushing), fainting or loss of consciousness, dizziness, and diarrhea. These are not all the possible side effects of TYVASO. Call your doctor for medical advice about side effects.

You may report side effects to the FDA at www.fda.gov/MedWatch or call 1-800-FDA-1088.

The risk information provided here is not comprehensive. To learn more about TYVASO, talk with your healthcare provider. Please see **Full Prescribing Information, Patient Product Information**, and the **TD-100** and **TD-300** TYVASO® Inhalation System Instructions for Use manuals at www.tyvaso.com or call 1-877-UNITHER (1-877-864-8437).

TYVISIconOct19



Breathe it in...

And get back to what really matters most



- ✓ TYVASO is inhaled directly to your lungs so it can quickly deliver medicine to the site of disease
- ✓ Reach your target dose quickly
- ✓ Walk farther and stay more active with fewer symptoms
- ✓ Take your medicine at home or on-the-go
- ✓ Receive support throughout your treatment journey

TYVASO DELIVERS

Join the TYVASO Support Program

If you are considering or are already taking TYVASO, sign up for the TYVASO Support Program to receive support with helpful tips and resources.

Sign up today at www.tyvaso.com or with the registration card in the pocket of this brochure.

Important Safety Information for TYVASO

Before you take TYVASO, tell your healthcare provider about all of your medical conditions, including if you (continued):

- Are pregnant or plan to become pregnant. It is not known if TYVASO will harm your unborn baby.
- Are breast-feeding or plan to breast-feed. It is not known if TYVASO passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with TYVASO.

Please see Important Safety Information throughout and on page 11 and accompanying Full Prescribing Information in pocket.

